

National Kidney Foundation
K/DOQI Guidelines: Kidney/Disease Outcome Quality Initiative*

Recommendations for Testing Frequency

Patients with Intact PTH (iPTH) within reference range for respective CKD Stage				
eGFR Value (ml/min/1.73m ²)	CKD Stage	Reference Range of iPTH (pg/mL)	Frequency of iPTH, Calcium and Phosphorus Testing	Frequency of Hydroxy Vitamin D Testing
30-59	3	35-70	Every 12 Months	N/A
15-29	4	70-110	Every 3 Months	N/A

Patients with Intact PTH (iPTH) above reference range for respective CKD Stage				
eGFR Value (ml/min/1.73m ²)	CKD Stage	Reference Range of iPTH (pg/mL)	Frequency of iPTH, Calcium and Phosphorus Testing	Frequency of Hydroxy Vitamin D Testing
30-59	3	35-70	Every 3 Months	Every 12 Months if >30 ng/mL Every 6 Months if <30 ng/mL
15-29	4	70-110	Every 3 Months	Every 12 Months if >30 ng/mL Every 6 Months if <30 ng/mL

*Clinical Practice Guidelines for CKD: Evaluation, Classification and Stratification.
American Journal of Kidney Disease, February 2002.